

Pre-Myelogram Medication Screening List

Amantadine (Symmetrel)
Aminophylline
Amitriptyline (Elavil)
Amitriptyline/chlordiazepoxide (Limbitrol DS)
Amitriptyline/perphenazine (Etrafon)
Amoxapine (Asendin)
Amphetamine
Amphetamine mixture (Adderall)
Aripiprazole (Abilify)
Asenapine (Saphris)
Bupropion (Wellbutrin, Zyban)
Bupirone (Buspar)
Citalopram (Celexa)
Chlorambucil (Leukeran)
Chloroquine (Aralen)
Chlorpromazine (Thorazine)
Clomipramine (Anafranil)
Clozapine (**Fazaclor, Versacloz**)
Cycloserine (Seromycin)
Cyclosporine (Sandimmune, Neoral)
Desipramine (Norpramin)
Desvenlafaxine (Pristiq)
Dexmethylphenidate (Focalin XR)
Dextroamphetamine (Dexedrine, etc)
Doxapram (Dopram)
Doxepin (Adapin, Sinequan)
Duloxetine (Cymbalta)
Droperidol
Ephedrine
Ertapenem (Invanz)
Escitalopram (Lexapro)
Fentanyl (Duragesic)
Fluoxetine (Prozac)
Fluphenazine (Prolixin, Permitil)
Fluvoxamine (Luvox)
Haloperidol (Haldol)
Hydroxychloroquine (Plaquenil)
Iloperidone (Fanapt)
Imipenem-cilastatin (Primaxin)
Imipramine (Tofranil)
Isocarboxazid (Marplan)
Isoniazid
Levomilnacipran (Fetzima)
Lindane
Linezolid (Zyvox)
Lisdexamfetamine (Vyvanse)
Lithium (Eskalith, Lithobid, Lithotabs)
Loxapine (Loxitane)
Lurasidone (Latuda)
Maprotiline (Ludiomil)
Mefloquine
Meperidine (Demerol)
Meropenem (Merrem)
Mesoridazine (Serentil)
Methamphetamine (Desoxyn)
Methylphenidate (Ritalin, **Concerta, Metadate, Methylin**)
Methylphenidate patch (Daytrana)
Metronidazole (Flagyl)
Milnacipran (Savella)
Mirtazapine (Remeron)
Molindone (Moban)
Nefazodone (Serzone)
Nortriptyline (Pamelor, Aventyl)
Olanzapine (Zyprexa)
Olanzapine/Fluoxetine (Symbyax)
Oseltamivir (Tamiflu)
Paliperidone (Invega)
Paroxetine (Paxil)
Pentazocine-naloxone
Perphenazine (Trilafon)
Phenelzine (Nardil)
Pimozide (Orap)
Procarbazine (Matulane)
Prochlorperazine (Compazine)
Promazine (Sparine)
Promethazine (Phenergan)
Protriptyline (Vivactil)
Pseudoephedrine
Quetiapine (Seroquel)
Rasagiline (Azilect)
Reserpine
Risperidone (Risperdal)
Selegiline (**Emsam, Zelapar**)
Sertraline (Zoloft)
Tapentadol (Nucynta)
Theophylline (**Theo-24, Elixophyllin**)
Thioridazine (Mellaril)
Thiothixene (Navane)
Tramadol (Ultram)
Tranlycypromine (Parnate)
Trazodone (Desyrel)
Trifluoperazine (Stelazine)
Triflupromazine (Vesprin)
Trimipramine (Surmontil)
Venlafaxine (Effexor)
Vilazodone (Viibryd)
Vortioxetine (Brintellix)
Ziprasidone (Geodon)

Drugs that lower the seizure threshold

1. Phenothiazines
2. Antipsychotics
3. Tricyclic antidepressants
4. CNS stimulants
5. Monoamine oxidase inhibitors
6. **Certain antibiotics**

Antiplatelet/Anticoagulant Agents[#]

DRUG	WHEN TO STOP
Abciximab (ReoPro [®])	48 hours
Apixaban (Eliquis [®])	24 to 48 hours*
Argatroban	2 hours
Aspirin	Does NOT need to be stopped
Bivalirudin (Angiomax [®])	2 to 6 hours
Cilostazol (Pletal [®])	48 hours
Clopidogrel (Plavix [®])	7 days
Dabigatran (Pradaxa [®])	1 to 6 days*
Dalteparin (Fragmin [®])	12 to 24 hours*
Dipyridamole (Persantine [®])	48 hours
Dipyridamole/aspirin (Aggrenox [®])	48 hours
Edoxaban (Savaysa [®])	12 to 24 hours*
Enoxaparin (Lovenox [®])	12 to 24 hours*
Eptifibatide (Integrilin [®])	4 to 8 hours
Fondaparinux (Arixtra [®])	24 hours*
Heparin, unfractionated	12 hours
NSAIDs	Do NOT need to be stopped
Prasugrel (Effient [®])	7 days
Rivaroxaban (Xarelto [®])	2 to 4 days*
Ticagrelor (Brilinta [®])	5 days
Ticlopidine (Ticlid [®])	7 days
Tinzaparin (Innohep [®])	12 to 24 hours
Tirofiban (Aggrastat [®])	8 hours
Vorapaxar (Zontivity [®])	4 weeks
Warfarin (Coumadin [®])	5 days or based on INR*

* Anticoagulant duration of effect depends on dose, age, renal function & concomitant medications; consult with pharmacy or prescribing information for further information.

[#] Do not stop therapy without consulting with patient's prescriber to assess risk vs benefit.