

One Step miRNA G0774 pFBAAVmU6mcs CMVmCherry SV40pA



Plasmid Features:


<u>Coordinates</u>	<u>Feature</u>
194-348	Tn7L
376-511	SV40pA (complementary-outside ITRs)
678-818	AAV2 ITR (141bp)
877-1188	EcoRI mU6 Promoter
1209-1298	MCS
1332-1337	Pol III Terminator EcoRI
1350-1929	CMV Promoter
1937-2665	mCherry
2718-2854	SV40pA
2926-3055	AAV2 ITR (130bp)
3612-4145	Gentamicin
4212-4436	Tn7R
5500-6360	B-lactamase (Ampicillin) Complementary

Antibiotic Resistance: Ampicillin and Gentamicin
Bacterial Backbone: pFastBac™ (Invitrogen)

Note: To check the integrity of the AAV ITR's perform single restriction enzyme digestions with XmaI, BssHII, and MscI.

This plasmid is designed for ease of cloning miRNA oligos directly into the pFastBac plasmid backbone.

The plasmid may also be used as a template for the mU6 promoter to tail on shRNA constructs. Constructs can then be cloned back into this plasmid with EcoRI sites to replace the original mU6 –pol III Terminator cassette. Or, alternatively, cloned into the G0451 or G0619 pFBAAVmcsCMVeGFP plasmids.



Red type indicates unique restriction site.

Multiple Cloning Site:

KpnI-PmeI-XhoI-SalI-ClaI-HindIII-EcoRV-EcoRI-XmaI-SmaI-BamHI-SpeI-XbaI-NotI.

The XhoI and XbaI sites recommended for miRNA cloning are unique. Methods for cloning miRNA can be found in this manual and are excerpts from *Generation of hairpin-based RNAi vectors for biological and therapeutic application.* Boudreau RL, Davidson BL. *Methods Enzymol.* 2012;507:233-251. doi:10.1016/B978-0-12-386509-00001-4

<http://www.sciencedirect.com/science/article/pii/B9780123865090000144>

>G0774 pFBAAVmU6mcsCMVmCherry 6946bp

CCATTCGCCATTCAGGCTGCAAATAAGCGTTGATATTCAGTCAATTACAAACATTAATAACGAAGAGATG
ACAGAAAAATTTTCATTCTGTGACAGAGAAAAAGTAGCCGAAGATGACGGTTTGTACATGGAGTTGGCA
GGATGTTTGATTAACAAACATAACAGGAAGAAAAATGCCCGCTGTGGGCGGACAAAATAGTTGGGAAC
GGGAGGGGTGGAATGGAGTTTTTAAGGATTATTTAGGGAAGAGTGACAAAATAGATGGGAACGGGTG
TAGCGTCGTAAGCTAATACGAAAATTAATAATGACAAAATAGTTTGGAACTAGATTTCACTTATCTGGTTC
GGATCTCCTAGGCGATATCAGTGATCAGATCCAGACATGATAAGATACATTGATGAGTTTGGACAAACCA
CAACTAGAATGCAGTGAAAAAATGCTTTATTTGTGAAATTTGTGATGCTATTGCTTTATTTGTAACCATTA
TAAGCTGCAATAAACAAGTTAACAACAACAATTGCATTCATTTTATGTTTCAGGTTCCAGGGGAGGTGTG
GGAGGTTTTTTAAAGCAAGTAAAACCTCTACAAATGTGGTATGGCTGATTATGATCCTCTAGTACTTCTCG
ACAAGCTTACATTATTGAAGCATTATCAGGGTTATTGTCTCAGACCTGCAGGCAGCTGCGCGCTCGCTC
GCTCACTGAGGCCGCCCGGGCAAAGCCCGGGCGTCCGGGCGACCTTTGGTCCCGGCCCTCAGTGAGCG
AGCGAGCGCGCAGAGAGGGAGTGGCCAACTCCATCACTAGGGGTTCTTGTAGTTAATGATTAACCCGC
CATGCTACTTATCTACGTAGCCATGCTCTAGTGAATTCGACGCCGCCATCTCTAGGCCCGCGCCGGCCCC
CTCGCACAGACTTGTGGGAGAAGCTCGGCTACTCCCCTGCCCGGTTAATTTGCATATAATATTTCTAG
TAACTATAGAGGCTTAATGTGCGATAAAAGACAGATAATCTGTTCTTTTTAATACTAGCTACATTTTACAT
GATAGGCTTGGATTTCTATAAGAGATACAAATACTAAATTATTATTTAAAAAACAGCACAAAAGGAACT
CACCTAACTGTAAGTAATTGTGTGTTTTGAGACTATAAATATCCCTTGGAGAAAAGCCTTGTTCGCTT
TAGTGAACCGTCAGATGGTACCGTTTAACTCGAGGTCGACGGTATCGATAAGCTTGATATCGAATTCCT
GCAGCCCGGGGATCCACTAGTTCTAGAGCGCCGCCACAGCGGGGAGATCCAGACATGATAAGATAC
ATTTTTGAATTCGGATCCTTATTAATAGTAATCAATTACGGGGTCATTAGTTCCATAGCCCATATATGGAGT
TCCGCGTTACATAACTTACGGTAAATGGCCCCCTGGCTGACCGCCCAACGACCCCGCCATTGACGT
CAATAATGACGTATGTTCCCATAGTAACGCCAATAGGGACTTTCCATTGACGTCAATGGGTGGAGTATTT
ACGGTAAACTGCCCACTTGGCAGTACATCAAGTGTATCATATGCCAAGTACGCCCCCTATTGACGTCAAT
GACGGTAAATGGCCGCCTGGCATTATGCCAGTACATGACCTTATGGGACTTTTCTACTTGGCAGTACA
TCTACGTATTAGTCATCGCTATTACCATGGTGATGCGGTTTTGGCAGTACATCAATGGGCGTGGATAGCG
GTTTGACTCACGGGGATTTCCAAGTCTCCACCCATTGACGTCAATGGGAGTTTGTGTTTGGCACCAAAT
CAACGGGACTTTCCAAAATGTCGTAACAACCTCCGCCCATTTGACGCAAAATGGGCGGTAGGCGTGTACGG
TGGGAGGTCTATAAAGCAGAGCTGGTTTGTGAAACCGTCAGATCCGCTAGCGCCACCATTGGTGAGCAA
GGGCGAGGAGGATAACATGGCCATCATCAAGGAGTTCATGCGCTTCAAGGTGCACATGGAGGGCTCCG
TGAACGGCCACGAGTTCGAGATCGAGGGCGAGGGCGAGGGCCGCCCTACGAGGGCACCCAGACCGC
CAAGCTGAAGGTGACCAAGGGTGGCCCCCTGCCCTTGCCTGGGACATCCTGTCCCCTCAGTTCATGTA
CGGCTCAAGGCCTACGTGAAGCACCCGCGGACATCCCCGACTACTTGAAGCTGTCTTCCCAGAGGG
CTTCAAGTGGGAGCGCGTGATGAACTTCGAGGACGGCGGCGTGGTGACCGTGACCCAGGACTCCTCCC
TGCAGGACGGCGAGTTCATCTACAAGGTGAAGCTGCGCGGCACCAACTTCCCCTCCGACGGCCCCGTAA
TGCAGAAGAAGACCATGGGCTGGGAGGCCTCCTCCGAGCGGATGTACCCCGAGGACGGCGCCCTGAAG
GGCAGATCAAGCAGAGGCTGAAGCTGAAGGACGGCGGCCACTACGACGCTGAGGTCAAGACCACCTA
CAAGGCCAAGAAGCCCGTGCAGCTGCCGGCGCCTACAACGTCAACATCAAGTTGGACATCACCTCCA
CAACGAGGACTACACCATCGTGGAAACAGTACGAACGCGCCGAGGGCCGCCACTCCACCGGCGGCATGG
ACGAGCTGTACAAGTAAAGACTCAATTTCGCCCTATAGTGAGTCGTATTACGCGCGCAGCGGCCGACCA
TGGCCCAACTTGTGTTATTGCAGCTTATAATGGTTACAAATAAAGCAATAGCATCACAAATTTACAAATAA
AGCATTTTTTTCACTGCATTCTAGTTGTGGTTTTGTCCAAACTCATCAATGTATCTTATCATGTCTGGATCTC
CGGACACGTGCGGACCGAAGTACAGCATGGCTACGTAGATAAGTAGCATGGCGGGTTAATCATTAACTA
CAAGGAACCCCTAGTGATGGAGTTGGCCACTCCCTCTGCGCGCTCGCTCGCTCACTGAGGCCGGGCG
ACCAAAGTTCGCCCGACGCCCGGGCGGCCCTCAGTGAGCGAGCGAGCGCGCAGCTGCCTGCAGGCATG

CAAGCTGTAGCCAACCACTAGAACTATAGCTAGAGTCCTGGGCGAACAAACGATGCTCGCCTTCCAGAA
AACCGAGGATGCGAACCACTTCATCCGGGGTTCAGCACCACCGGCAAGCGCCGCGACGGCCGAGGTCTT
CCGATCTCCTGAAGCCAGGGCAGATCCGTGCACAGCACCTTGCCGTAGAAGAACAGCAAGGCCGCCAA
TGCCTGACGATGCGTGGAGACCGAAACCTTGCCTCGTTCGCCAGCCAGGACAGAAATGCCTCGACTTC
GCTGCTGCCCAAGGTTGCCGGGTGACGCACACCGTGGAAACGGATGAAGGCACGAACCCAGTTGACAT
AAGCCTGTTGCGTTCGTAACCTGTAATGCAAGTAGCGTATGCGCTCACGCAACTGGTCCAGAACCTTGAC
CGAACGCAGCGGTGGTAACGGCGCAGTGGCGGTTTTTCATGGCTTGTATGACTGTTTTTTTTGTACAGTCT
ATGCCTCGGGCATCCAAGCAGCAAGCGCTTACGCCGTGGGTGATGTTTGTATGACTGTTTTTTTTGTACAGTCT
GATGTTACGCAGCAGCAACGATGTTACGCAGCAGGGCAGTCGCCCTAAAACAAAGTTAGGTGGCTCAAG
TATGGGCATCATTGCGACATGTAGGCTCGGCCCTGACCAAGTCAAATCCATGCGGGCTGCTCTTGATCTT
TTCGGTTCGTGAGTTCGGAGACGTAGCCACCTACTCCAACATCAGCCGGACTCCGATTACCTCGGGAAC
TTGCTCCGTAGTAAGACATTCATCGCGCTTGCTGCCTTCGACCAAGAAGCGGTTGTTGGCGCTCTCGCGG
CTTACGTTCTGCCAAGTTTGAGCAGCCGCGTAGTGAGATCTATATCTATGATCTCGCAGTCTCCGGCGA
GCACCGGAGGCAGGGCATTGCCACCGCGCTCATCAATCTCCTCAAGCATGAGGCCAACGCGCTTGGTG
CTTATGTGATCTACGTGCAAGCAGATTACGGTGACGATCCCGCAGTGGCTCTCTATAAAAGTTGGGCAT
ACGGGAAGAAGTGATGCACTTTGATATCGACCCAAGTACCGCCACCTAACAATTCGTTCAAGCCGAGAT
CGGCTTCCCGGCCGCGGAGTTGTTGCGTAAATTGTCACAACGCCGCGAATATAGTCTTTACCATGCCCTT
GGCCACGCCCTCTTAATACGACGGGCAATTTGCACCTTCAGAAAATGAAGAGTTTGCTTTAGCCATAAC
AAAAGTCCAGTATGCTTTTTACAGCATAAAGTGGACTGATTTGAGTTTACAACCTATTCTGTCTAGTTAAG
ACTTTATTGTCATAGTTTAGATCTATTTTTGTTGAGTTTAAAGCTTTATTGTCGGCCACACCCGCTTACGCA
GGGCATCCATTTACTCAACCGTAACCGATTTTGCAGGTTACGCGGCTGGTCTATGCGGTGTGAAT
ACCGCACAGATGCGTAAGGAGAAAATACCGCATCAGGCGCTTTCGCTTCCGCTCCTCGCTCACTGACTCGCT
GCGCTCGGTGTTGCGCTGCGGCGAGCGGTATCAGCTCACTCAAAGGCGGTAATACGTTTATCCACAGA
ATCAGGGGATAACGCAGGAAAGAACATGTGAGCAAAGGCCAGCAAAGGCCAGGAACCGTAAAAAG
GCCGCGTTGCTGGCGTTTTTCCATAGGCTCCGCCCCCTGACGAGCATCACAAAATCGACGCTCAAGT
CAGAGGTGGCGAAACCCGACAGGACTATAAAGATACCAGGCGTTTCCCCCTGGAAGCTCCCTCGTGCGC
TCTCCTGTTCCGACCCTGCCGCTTACCGGATACCTGTCCGCTTTCTCCCTTCGGGAAGCGTGGCGCTTT
CTCATAGCTCACGCTGTAGGTATCTCAGTTCGGTGTAGGTCGTTGCTCCAAGCTGGGCTGTGTGCACGA
ACCCCCCGTTACGCCGACCGCTGCGCCTTATCCGGTAACTATCGTCTTGAGTCCAACCCGTAAGACA
CGACTTATCGCCACTGGCAGCAGCCACTGGTAACAGGATTAGCAGAGCGAGGTATGTAGGCGGTGCTAC
AGAGTTCTTGAAGTGGTGGCCTAACTACGGCTACACTAGAAGAACAGTATTTGGTATCTGCGCTCTGCTG
AAGCCAGTTACCTTCGGA AAAAGAGTTGGTAGCTCTTGATCCGGCAAACAACACCGCTGGTAGCGGT
GGTTTTTTTTGTTTGAAGCAGCAGATTACGCGCAGAAAAAAGGATCTCAAGAAGATCCTTTGATCTTTT
TACGGGGTCTGACGCTCAGTGAACGAAAACACTCACGTTAAGGGATTTGGTCATGAGATTATCAAAAAGG
ATCTTACCTAGATCCTTTTAAATTA AAAATGAAGTTTTAAATCAATCTAAAGTATATATGAGTAAACTTGG
TCTGACAGTTACCAATGCTTAATCAGTGAGGCACCTATCTCAGCGATCTGTCTATTTTCGTTTATCCATAGT
TGCTGACTCCCGTTCGTGTAGATAACTACGATACGGGAGGGCTTACCATCTGGCCCCAGTGTGCAAT
GATACCGCGAGACCCACGCTCACCGGCTCCAGATTTATCAGCAATAAACCAGCCAGCCGGAAGGGCCG
AGCGCAGAAGTGGTCTGCACTTTATCCGCCTCCATCCAGTCTATTAATTGTTGCCGGGAAGCTAGAGT
AAGTAGTTCGCCAGTTAATAGTTTGCGCAACGTTGTTGCCATTGCTACAGGCATCGTGGTGTACGCTCG
TCGTTTGGTATGGCTTCATTCAGCTCCGTTCCCAACGATCAAGGCGAGTTACATGATCCCCATGTTGT
GCAAAAAGCGGTTAGCTCCTTCGGTCTCCGATCGTTGTGAGAAGTAAGTTGGCCGAGTGTATCACT
CATGGTTATGGCAGCACTGCATAATTCTCTTACTGTATGCCATCCGTAAGATGCTTTTTCTGTGACTGGTG
AGTACTCAACCAAGTCATTCTGAGAATAGTGTATGCGGCGACCGAGTTGCTCTTGCCCGCGTCAATACG
GGATAATACCGCGCCACATAGCAGA ACTTTAAAAGTGCTCATCATTGGAAAACGTTCTTTCGGGGCGAAA
ACTCTCAAGGATCTTACCGCTGTTGAGATCCAGTTGATGTAACCCACTCGTGACACCAACTGATCTTCA
GCATCTTTTACTTTACCAAGCGTTTTCTGGGTGAGCAAAAACAGGAAGGCAAATGCCGCAAAAAGGGA
ATAAGGGCGACACGGAAATGTTGAATACTCATACTCTTCTTTTTCAATATTATTGAAGCATTATCAGGG
TTATTGTCTCATGAGCGGATACATATTTGAATGTATTTAGAAAAATAAACAATAGGGGTTCCGCGCACAT
TTCCCGAAAAGTGCCACCTAAATTGTAAGCGTTAATATTTTGTAAAATTCGCGTTAAATTTTTGTTAAAT
CAGCTCATTTTTTAACCAATAGGCCGAAATCGGCAAAATCCCTTATAAATCAAAGAATAGACCGAGATA
GGGTTGAGTGTGTTCCAGTTTGAACAAGAGTCCACTATTAAGAACGTGGACTCCAACGTCAAAGGGC
GAAAAACCGTCTATCAGGGCGATGGCCCACTACGTGAACCATCACCTAATCAAGTTTTTTGGGGTCGA
GGTGCCGTA AAGCACTAAATCGGAACCTAAAGGGAGCCCCGATTTAGAGCTTGACGGGGAAAGCCG
GCGAACGTGGCGAGAAAGGAAGGGAAGAAAGCGAAAGGAGCGGGCGCTAGGGCGCTGGCAAGTGTAG
CGGTCACGCTGCGCGTAACCACCACACCCGCCGCGCTTAATGCGCCGCTACAGGGCGCGTC

Plasmid Quality Control ITR Integrity Digest:

Expected Fragments:


BssHII:	4600, 1918, 269, 85, 74
MscI:	3503, 1288, 1185, 970
XmaI:	4652, 1744, 539, 11
XbaI:	6946 (necessary restriction site for miRNA protocol)
XhoI:	6946 (necessary restriction site for miRNA protocol)

G0774 pFBAAVmU6mcsCMVmCherry SV40pA

Aldevron lot: 59163 2.3ug/ul 4-30-15


5-6-15 AN, 300ng/well

1kb+ Uncut XhoI BssHII MscI XmaI




Plasmid Quality Control Transfection:

Transfection in 293FT Cells 0.7E+06 c/well.
24hrs Post Transfection



G0774 pFBAAVmU6mcsCMVmCherry
1ug/well



G0760 pFBAAVmU6mcsCMVeGFP
1ug/well

Specific Information for cloning miRNA in the G0760 pFBAAVmU6mcsCMVeGFP or G0774 pFBAAVmU6mcsCMVmCherry One Step miRNA Plasmids:

Background:

The restriction sites of these miRNA Cloning vectors have been reconfigured to allow the direct cloning of miRNA annealed oligo PCR products in one round as opposed to the original method outlined in the references below through TopoBlunt:mU6 and then into the pFBAAV vector. One Step miRNA plasmids also allow for screening miRNA with eGFP or mCherry reporters not present in the TopoBlunt:mU6 vector. The mU6 promoter may be used as a template for existing shRNA strategies outlined by Boudreau et. al. Please refer to the original publications for working with shRNA constructs.

References:

Generation of hairpin-based RNAi vectors for biological and therapeutic application. Boudreau RL, Davidson BL. *Gene*. 2012;507:275-285. doi:10.1016/j.gene.2012.07.015. Epub 2012 Jul 15. <http://www.sciencedirect.com/science/article/pii/S014527511223865090000144>

siSPOTR: a tool for designing highly specific and potent siRNAs for human and mouse. Boudreau RL, Spengler RM, Hylock RH, Kusenda BJ, Davis HA, Eichmann DA, Davidson BL. *Nucleic Acids Res*. 2011 Dec 19;39(24):8711-8721. doi:10.1093/nar/nbr847. Epub 2011 Sep 27.

siSPOTR (siRNA Sequence Probability-of-Off-Targeting Reduction) <https://sispotr.icts.uiowa.edu/sispotr/index.html;jsessionid=E8EB7B91C5640FD84F7E72DC758505DD>

Rational design of therapeutic siRNAs: minimizing off-targeting potential to improve the safety of RNAi therapy for Huntington's disease. Boudreau RL, Spengler RM, Davidson BL. *Mol Ther*. 2011 Dec;19(12):2169-2179. doi:10.1038/mt.2011.185. Epub 2011 Sep 27.

Protocol:

Below are revised excerpts from **Generation of hairpin-based RNAi vectors for biological and therapeutic application** for miRNA design and cloning. Section numbers refer to the numbers in the original publication.

GENERATION OF HAIRPIN-BASED RNAi VECTORS FOR BIOLOGICAL AND THERAPEUTIC APPLICATION

Ryan L. Boudreau and Beverly Davidson

Contents

2.	Selecting Candidate siRNA Sequences	6
2.1.	Retrieve sequences for your desired target transcript(s)	6
2.2.	Select 22-nt long target sites	7
4.	Cloning Artificial miRNA Expression Vectors	7
4.1.	Considerations for designing artificial miRNAs	7
4.2.	Oligo design and cloning protocol for artificial miRNAs	8

2. Selecting Candidate siRNA Sequences

2.1. Retrieve sequences for your desired target transcript(s)

Choose your target gene(s) of interest and retrieve the relevant mRNA sequence(s) from NCBI, Ensembl, UCSC Genome Browser, or other available databases. We leave it to the reader's responsibility to further investigate target transcripts for variants which may result from RNA processing events (e.g., alternative splicing, alternative polyadenylation, and RNA editing, among others) that may influence the ability to target certain regions within the transcript. In general, we target the coding region; however, targeting the 5'- and 3'-UTR sequences is possible. Careful consideration for the target sequence

with regard to the project objectives is important before proceeding with design and screening of inhibitory RNA sequences. For instance, the reader should consider whether allele- or splice-isoform-specific silencing is desirable or whether sequence conservation is important, allowing the RNAi vectors to be tested in multiple species.

2.2. Select 22-nt long target sites

Identifying potent and highly specific siRNA sequences is not trivial. Numerous empirical evaluations of large-scale siRNA knockdown data have allowed researchers to establish several siRNA design guidelines (Khvorova et al., 2003; Matveeva et al., 2007). For example, one key consideration is that siRNA sequences be selected or manipulated to promote accurate loading of the antisense guide strand into RISC, leaving the sense strand to be degraded (Leuschner et al., 2006; Matranga et al., 2005). Furthermore, GC-content and positional nucleotide preferences also influence siRNA efficacy. Given the multifaceted nature of designing optimal siRNAs, we direct the reader to additional literature on the subject (Birmingham et al., 2007; Davidson and McCray, 2011; Jackson and Linsley, 2010). Also, there are numerous publicly available siRNA design tools online. It is important to note that siRNA design rules serve more as guidelines, and that sequences adhering to them may not silence and vice versa.

NEW: We have developed an online tool, siSPOTR (siRNA Seed Probability of Off-Target Reduction) to assist users with designing siRNAs with low off-targeting potential (https://research.icts.uiowa.edu/sispotr_review2011/)¹. siSPOTR searches user-defined target sequences for siRNAs that pass strand-biasing and GC% filters and outputs the candidates with the lowest POTS values. The output provides detailed off-target information for each siRNA including i) the number of 3'-UTRs containing each seed site type, ii) the putative off-target transcripts, and iii) counts of each seed site type on a per transcript basis. siSPOTR also determines if the siRNA seed matches that of a known miRNA. Recognizing the ease of purchasing pre-validated siRNAs, an additional tool allows users to input siRNA sequences to obtain relevant off-targeting scores and information. <https://sispotr.icts.uiowa.edu/sispotr/index.html;jsessionid=E8EB7B91C5640FD84F7E72DC758505DD>

To date, no algorithm guarantees silencing efficacy, and it is most recommend that the user pick three to four candidates for screening. Some investigators screen 10-20 candidates at a time. Tips and short cuts for working with multiple candidates are outlined in the cloning information below.

Here, we will describe a basic strategy for siRNA target site selection which incorporates the most important criteria for promoting efficacy, in addition to certain rules that are specific to the design of hairpin-based RNAi expression vectors.

We identify 22-nt target sites within the target transcript that adhere to four criteria: (1) high propensity to primarily load the antisense guide strand into RISC, (2) GC content between 20% and 70%, (3) void of restriction enzyme sites relevant to downstream applications (e.g., cloning RNAi expression cassettes into viral vector systems), (4) lacking a stretch of four continuous A or T nucleotides (i.e., AAAA or TTTT). The latter prevents premature transcription termination from Pol III promoters which typically terminate at stretches of four to six T's. Strand biasing is determined by the thermodynamic stabilities present at the ends of the siRNA duplex. To achieve faithful loading of the antisense strand, the duplex must be designed such that there is strong G–C base pairing present at the 5'-end of the sense (passenger) strand and weak A/G–U base pairing at the opposing terminus (Khvorova et al., 2003; Schwarz et al., 2003). The RISC complex selects the strand with the weakest 5'-end thermodynamic stability, in this case, the antisense strand (Fig. 14.1). Hence, we select target sequences that have G or C nucleotides at positions 3 and 4 (note: positions 1 and 2 correspond to the dinucleotide 3'-overhang of the antisense strand) and A, T, or C nucleotides at positions 20, 21, and 22. The C is allowed at the latter positions because we can destabilize the resulting G–C base pairs by converting them to G–U base pairs (G and U weakly pair in RNA) without altering the antisense strand sequence. Next, the GC content of the 22-nt site is calculated by dividing the number of G or C nucleotides by 22. For instance, the target site shown in Fig. 14.1 has 12 G or C nucleotides, resulting in a GC content of 55% which is within the acceptable range. Finally, we avoid stretches of A's and T's and relevant restriction enzyme sites for reasons mentioned above. Note: the user must also avoid creating these elements upon incorporating the siRNA into shRNA or artificial miRNA scaffolds.

4. Cloning Artificial miRNA Expression Vectors

4.1. Considerations for designing artificial miRNAs


Artificial miRNAs, or miRNA shuttles, are designed to mimic naturally occurring pri-miRNAs, for which the Drosha-DGCR8 and Dicer cleavage sites have been mapped and experimentally validated. With this information, the identity of the small RNA duplex which is processed from the initial stem-loop transcript is known. For artificial miRNAs, this region is replaced by siRNA duplexes, thus creating a miRNA-based hairpin which serves to shuttle siRNA sequences into the RNAi pathway. An important consideration for designing artificial miRNAs is to maintain the structural and sequence recognition motifs required for appropriate processing. Drosha-DGCR8 binds to regions of single-stranded nature located at the base of the pri-miRNA stem loop (Han et al., 2004; Zeng and Cullen, 2005). Thus, including 50–100 nts of the flanking sequences (5' and 3') native to the pri-miRNA will help to ensure that the stem-loop base folds properly to promote cleavage at the intended site. Artificial miRNAs have been generated using a number of naturally occurring pri-miRNAs as scaffolds for siRNA sequences (Chung et al., 2006; Tsou et al., 2011; Zeng et al., 2002). Here, we will describe a method to generate artificial miRNAs based on the natural human miR-30 pri-miRNA transcript. For ease of cloning, we only include a minimal amount of natural miR-30 flanking sequences, while additional 5'- and 3'-sequences are derived from the mouse U6 expression vector. We have characterized these vectors for appropriate expression and processing using small transcript northern blot, small RNA RT-PCR and RACE analyses (Boudreau et al., 2008; Chen et al., 2005). We recommend that newly designed miRNA shuttles

be similarly defined with these techniques, since simply switching promoters, restriction enzyme sites, or expression contexts (e.g., embedding the miRNA-based stem loop in the 3'-UTR or intron of a reporter gene) can alter the pri-miRNA structure and subsequent processing.

4.2. Oligo design and cloning protocol for artificial miRNAs

The following steps are outlined in Fig. 14.3.


Step 1. Start with artificial miRNA backbone sequence.



Step 2. Insert sense and antisense sequences.




Step 3. Manipulate 5' bulge nucleotide to induce a mismatch.



Step 4. Convert C's to T's at sense 3' end to impart duplex instability.



Step 5. Convert all T's to U's and fold RNA using UNAFold



Step 6. Design and order DNA oligos.



Step 7. Anneal and polymerase extend oligos.



Step 8. PCR purification, digestion and cloning to pTOPObluntII-U6



Figure 14.3. Artificial miRNA cloning scheme. The details for designing artificial miRNAs and cloning them into the mU6 expression vector are described in the text (Section 4.2, Steps 1–8). Shown here is the example of incorporating the eGFP-targeted siRNA into an artificial miRNA scaffold.

How to use the siSPOTR tool:

- Input sequence(s) for siRNA design at <https://sispotr.icts.uiowa.edu/sispotr/tools/sispotr/design.html> Click the appropriate button to paste or upload the sequence(s). Single sequences can be in plain or FASTA format. Multiple sequences require FASTA formatting.
- Select the number of candidates desired in the output.

Sigma-Aldrich, or Invitrogen) using the smallest synthesis scale available and standard desalting purification. Upon receipt, the oligos can be reconstituted in purified water to make a 100- μ M stock.

Hairpin 86mer

CTCGAGTGAGCGcTGCAGCCAGGACTGCGCTAAaCTGTAAAGCCACAGATGGGcTTAGCGCAGTCTGGCTGcATCGCCTACTAGT

Hairpin-broken into two primers to be annealed and extended: (Random A's added to the ends for digest)

Oligo 1 miRNA For: 5' AAAACTCGAGTGAGCGcTGCAGCCAGGACTGCGCTAAaCTGTAAAGCCACAGATGGG 3' 58mer

miRNA Rev: 3' CTGTAAAGCCACAGATGGGcTTAGCGCAGTCTGGCTGcATCGCCTACTAGT 5' 57mer

Oligo 2 (rev and comp:) 5' AAAAAGTAGTAGGCGATGCAGCCAGGACTGCGCTAAaCCCATCTGTGGCTTTACAG 3'

Step 7.

Anneal and polymerase extend Oligo 1 and Oligo 2.

Reaction conditions

5 μ l 10 \times reaction buffer with MgCl₂

1 μ l Oligo 1 (100 μ M)

1 μ l Oligo 2 (100 μ M)

1 μ l dNTPs (10 mM)

0.5 μ l enzyme mix (i.e., polymerase)

41.5 μ l dH₂O

Incubate the reaction in a thermocycler:

94 $^{\circ}$ C for 2 min (denaturation)

54 $^{\circ}$ C for 1 min (annealing)

72 $^{\circ}$ C for 15 min (extension)

Step 8.

PCR purification, digestion, and clone directly into G0760 pFBAAVmU6mcsCMVeGFP or G0774

pFBAAVmU6mcsCMVmCherry.

Step 8.1.

Purify the extended product using a PCR Purification kit (QIAquick; Qiagen or other) as per the manufacturer's instructions and elute in 20-30 μ l water.

Timesaving Tips for working with 15-20 candidates: Anneal and extend the oligos pairs individually as above. Run a sample on a gel to ensure that the extension worked for each pair and that they are the proper size and roughly the same molar ratio. Pool 5ul of each confirmed double stranded hairpin pair and proceed to digest, and ligate the product as if it is one sample. This reduces the sample handling greatly for several steps. Pick 30-48 minipreps from the resulting ligation. Process minipreps and sequence. At the University of Iowa, setting up sequencing reactions in a plate format with multiple samples greatly reduces the cost. Screen sequences for all of the original intended miRNA hairpin sequences. Typically, the return is 95-100%.

Step 8.2.

Digest the product with XhoI and SpeI by adding the following to the 30 μ l eluate:

4 μ l 10 \times restriction enzyme buffer (NEB2; New England Biolabs)

4 μ l 10 \times BSA (10 mg/ml)

1 μ l SpeI (10 units/ μ l)

1 μ l XhoI (20 units/ μ l)

Incubate reaction at 37 $^{\circ}$ C for 4 h to overnight.

Step 8.3.

Digest 3-4 μ g of the G0760 pFBAAVmU6mcsCMVeGFP or G0774 pFBAAVmU6mcsCMVmCherry expression plasmid with XhoI and XbaI. This plasmid contains the mouse U6 promoter followed by a multiple cloning site (MCS) and a Pol III termination signal (six T's) (Fig. 14.4). XbaI- (vector) and SpeI (artificial miRNA)-cleaved sites produce compatible sticky ends for ligation. Although there is a SpeI site in the G0760 pFBAAVmU6mcsCMVeGFP and G0774

pFBAAVmU6mcsCMVmCherry multiple cloning sites, we have found in prior studies that ligation to the XbaI site produces a stem loop which is more efficiently processed, yielding higher antisense RNA levels and more potent gene silencing (Boudreau et al., 2008).

Step 8.4.

Gel purify the digested fragments. We typically run the digested artificial miRNA inserts on a 2% agarose gel and excise the ~100-bp band. The digested pFBAAVmU6mcsCMVeGFP (or mCherry) is run on a 1% agarose gel and the ~7-kb fragment is excised. Gel extraction can be performed by various means.

Step 8.5.

Perform ligation and bacterial transformation using standard protocols. We ligate 6 ng of insert to 50 ng of vector and incubate at room temperature for 1 h before transformation. The pFBAAV plasmid backbone is both ampicillin and

gentamicin resistant, and thus, transformed bacteria should be grown on LB agar plates containing a final concentration of 100ug/ml ampicillin and 7ug/ml gentamicin. We highly recommend the use of both antibiotics. The gentamicin resistance is contained within the transposable elements. The ampicillin resistance is outside the transposable elements. The possibility for recombination at the transposable elements or ITRs exists anytime you amplify the plasmid. Recombination at the transposable elements may lead to amp resistant false positives that contain no gene of interest on ampicillin only plates. Because the pFBAAVmU6mcsCMVeGFP vector is digested with two non-cohesive enzymes (XhoI and XbaI), the likelihood of intramolecular vector ligation is minimal, and few, if any, A/G-resistant colonies grow on “vector-only” control plates. However, we recommend the vector be treated with alkaline phosphatase after restriction enzyme digestion to further minimize the potential for background colonies. We typically pick four to six colonies per artificial miRNA construct and grow each of them overnight in 3 ml liquid LB cultures containing 100ug/ml ampicillin and 7ug/ml gentamicin. The following day, minipreps are performed using a miniprep kit which yields plasmid DNA of sufficient quality for cell culture applications.

Step 8.6.

Screen for positive clones by EcoRI digestion and gel electrophoresis. Note: cloning an artificial miRNA into the XhoI and XbaI sites removes the internal EcoRI (in the MCS), and thus, positive clones can be screened for by EcoRI digestion. pFBAAVmU6mcsCMVeGFP or mCherry vector-only plasmid will yield 6489bp, 383bp, and 83bp fragments indicating a negative clone; we recommend digesting the parental pFBAAVmU6 vector alongside the potential artificial miRNA clones to serve as a reference. The positive constructs with successful insertion of an artificial miRNA will yield 6489bp and 483bp bands.

Step 8.7.

Sequence positive clones using the mU6 forward primer or CMVR primer. Although the orientation of the mU6miRNA cassette is independent of the CMVeGFP or CMVmCherry and can be in either orientation, we generally choose those in the forward orientation for ease of sequence and map preparation. Many people make mistakes when putting together sequences and maps of inserts in the reverse orientation from the body of the plasmid. The NotI site in the cassette in conjunction with an enzyme outside the cassette such as XmaI can be used to conveniently determine orientation. For example, a double digest with NotI and XmaI will yield expected fragments of 4652bp, 1728bp, 581bp, and 11bp with the cassette in the “Forward” orientation and 4652bp, 2119bp, 190bp, and 11bp with the cassette in the “Reverse” orientation.

Suggested Sequencing Primers:

mU6-F	ACAGACTTGTGGGAGAAGC
CMV-R	GTAGGAAAGTCCCATAAGGTCA

4.3. Materials for artificial miRNA cloning

- G0760 pFBAAVmU6mcsCMVeGFP plasmid
- G0774 pFBAAVmU6mcsCMVmCherry plasmid
- Overlapping DNA oligos for artificial miRNA
- Expand high-Fidelity Polymerase and buffers (Roche)
- Restriction enzymes and buffers (XhoI, SpeI, XbaI, EcoRI, NotI, XmaI, BssHII, and MscI)
- Gel electrophoresis equipment and reagents
- Gel extraction kit
- DNA ligase and buffer
- Chemical competent DH5a and Stbl3 bacterial cells
- Ampicillin and Gentamicin-containing LB agar growth plates
- Miniprep kit

Information and Cloning Suggestions for Working with pFBAAV Plasmids:

Characteristics of AAV:

Adeno-Associated virus (AAV) is a non-pathogenic helper dependant parvovirus. This is one of the most promising vehicles for gene delivery. Recombinant AAV vectors have predominantly episomal gene expression. AAV has long term expression in terminally differentiated cells. It can infect a wide range of cells, including non-dividing cells.

Insert Size:

The major disadvantage of the AAV vectors is the small genome, which limits the size of the transgene (from ITR to ITR) to about 4.7 Kb. Please consider the size of the promoter, polyA, any reporters, and your gene of interest when planning the cloning into an AAV shuttle vector. When the length of inserted DNA between the 2 ITRs is close the maximal allowed, i.e., 4.7Kb, the packaging efficiency decreases significantly.

AAV Production:

The University of Iowa Viral Vector Core uses two systems for AAV production: A Triple Transfection System and the Baculovirus System.

The Triple Transfection System may be used with any pAAV plasmid including the pFBAAV plasmids. A pHelper plasmid isolated from adenovirus, a Rep/Cap plasmid, and the AAV ITR plasmid package containing the gene of interest are transfected into 40 x 150mm plates of adherent 293FT cells per prep. This system is more labor intensive for production and purification but can be completed within three to four weeks of receiving the plasmid.

The services provided for production are:

- Transfection.
- Large-scale amplification of AAV vectors.
- The AAV vector is purified using either CsCl or an iodixanol gradient followed by ion-exchange or affinity chromatography.
- A physical titer in viral genomes per ml (vg/ml) is assessed by QPCR.
- **Sample required:** 1500ug of high quality AAV plasmid expressing the gene of interest at a concentration greater than 0.25ug/ul per prep. One disadvantage to the investigator in the triple transfection system is having to provide 1500ug of DNA for each subsequent prep and the possibility of recombination during the maxi prep process. Each maxi prep should be digested and evaluated for integrity.

The Baculovirus System can only be used with pFBAAV plasmids with Tn7 transposable elements to incorporate the ITRs and gene of interest into the Bacmid DNA. *Please note: not all AAV plasmids contain the Baculo elements so this system may not be an option for everyone. This system also relies on the gentamicin selection marker within the transposable elements. The system was developed by Dr. Robert Kotin at the National Institute of Health. The system is described in the following publication: "*Insect Cells as a Factory to Produce Adeno-Associated Virus Type 2 Vectors*" [Human Gene Therapy 13:1935-1943](#) (November 1, 2002). The Baculovirus System takes 5-6 weeks for production but has many advantages. The Baculovirus System can tolerate more toxic proteins driven by mammalian promoters as these promoters will not be active in the insect cells. This system is also generally more robust with higher titers, cost-effective, and less labor intensive for production. Once we have a Baculovirus stock of the investigator package, we can make additional preps readily and we have tested the Baculovirus stock to be stable for years.

The services provided for production are:

- Transposition of the gene interest into the DH10Bac bacmid DNA. Everything in the pFBAAV plasmid between the transposable elements (Tn7, ITR, Promoter, gene of interest, pA, ITR, gentamicin resistance, Tn7) is incorporated into the large 135,000kb bacmid DNA. Your plasmid must be gentamicin resistant for the transposition to work.
- Growth and midi prep preparation of Bacmid DNA.
- Transfection of the Bacmid DNA with gene of interest into Sf9 cells. Approximately 12mls of the P1 stage Baculovirus is harvested 5 days later.
- Titer of the Baculovirus P1 stock. We are able to make 1-3 preps of final AAV virus using the P1 stock depending on the titer of the stock and the desired MOI of infection for the final AAV prep.

- For multiple AAV virus preps the P1 stock is amplified to a larger volume P2 Baculovirus stock and titered.
- The final AAV virus is prepared in bioreactor bags on a rocker platform by dual infection with the package Baculovirus and the desired Rep/Cap Baculovirus into Sf9 cells. The Baculovirus provides pHelper function. The culture is harvested 3 days later.
- The AAV vector is purified using either CsCl or an iodixanol gradient followed by ion-exchange or affinity chromatography.
- A physical titer in viral genomes per ml (vg/ml) is assessed by QPCR.
- **Sample required:** 10ug of pFBAAV plasmid expressing the gene of interest at a concentration greater than 0.25ug/ul.

Bacterial Backbone:

All of our internal AAV plasmids contain Baculo compatible elements from Invitrogen pFastBac™. For information on the pFastBac backbone and the Baculovirus System, please review the Invitrogen manual: http://tools.invitrogen.com/content/sfs/manuals/bactobac_topo_exp_system_man.pdf

**Please note: not all AAV plasmids are compatible with the Baculo system.

ITRs (Inverted Terminal Repeats):

The ITRs for AAV were engineered into the Invitrogen pFastBac backbone by Dr. Robert Kotin.

Orientation and Updates:

The MCS and BgHpA were inserted in the antisense direction relative to the Invitrogen backbone and ITRs. Please note that the backbone and ITRs were re-sequenced and updated in 2012 and there are several changes. Please discard any older sequences. The nomenclature of 5'ITR (130bp) and 3'ITR (141bp) have been retained on some maps to identify the sequences. The orientation of the ITRs makes no difference to the formation of the viral vector.

Recombination:

Recombination is a possibility at both the transposable elements (Tn7L and Tn7R) and the Inverted Terminal Repeats (ITRs) in the pFBAAV plasmid. In order to check for and reduce the chance of recombination, we have the following recommendations.

Antibiotic Resistance:

The gentamicin resistant gene lies within the transposable elements and may be lost if the transposable elements recombine. We suggest using **both ampicillin and gentamicin** selection on plates and in your liquid media throughout the cloning process and during amplification to midi or maxi preps.

Ampicillin concentration: 100ug/ml final

Gentamicin Sulfate concentration: 7ug/ml final

E. Coli Competent Cell Recommendations:

We recommend using a stable E. coli strain such as **SURE2, Stable2, or Stable3** for transformation of your final plasmid product. These competent cell strains have been engineered to stop unwanted rearrangement events and lack the components of the pathways that catalyze the rearrangement and deletion of nonstandard secondary and tertiary structures, including cruciforms (caused by inverted repeats) and Z-DNA, that occur frequently in eukaryotic cells. Cloning, however, can be difficult in these strains. We recommend DH5a competent cells for sub-cloning.

Quality Control ITR Digest:

We require a quality control digest of the plasmid upon submission to ensure that there is no recombination before vector production is started. The ITRs in the pFBAAV plasmids have several convenient restriction sites to determine whether the ITRs are intact without sequencing. Sequencing the ITRs can be difficult due to their hairpin secondary structure. We suggest a single digest each of **BssHII, MscI, and XmaI**. It is a good idea to also run un-cut and linearized plasmid next to these digests. Determine your expected fragment sizes for each digest and check carefully to see that you get what you expect and have no extraneous bands. Be sure to check your final midi or maxi product (not just your miniprep) as recombination is possible during any amplification process.

Sequencing pFBAAV plasmids and ITRs

The majority of the plasmid and insert can be sequenced normally. Sequencing of the ITRs is difficult but possible. Sequencing of the ITRs or other difficult secondary structures may be aided by requesting "Extended Denaturation" or by requesting "Betaine Chemistry for AAV ITRs" in the comments section at the University of Iowa Sequencing Core. Non-UI investigators should consult with their own sequencing facility for help if sequencing of the ITRs is desired. We routinely request "Extended Denaturation" for all AAV plasmid sequencing.

There is no one set of primers that will always sequence through the ITRs. The ITR primers listed below have been used with success in many cases. Try sequencing in both directions: from the plasmid in and from your insert out.

Suggested Sequencing Primers for a variety of pFBAAV plasmids:

Below is a general list of commonly used primers at the University of Iowa Viral Vector Core. Please check to make sure the primers work for the shuttle plasmid you have received. There are many free primer design programs available to design primers that will sequence your insert.

BgHpA For	TGAGGAAATTGCATCGCATTGTCT
BgHpA Rev	AGGAAAGGACAGTGGGAGTG
pFBAAV (130bp ITR)	GCCTTGCTGTTCTTCTACGG
pFB587F (141bp ITR)	CTCTACAAATGTGGTATGGCTG
pFB214F (141bp ITR)	GGGGTGGAAATGGAGTTT
pFB158F (141bp ITR)	CATAACAGGAAGAAAATGCCCCG
EGFP-141Rev	GAACCTCAGGGTCAGCTTGCCGTA
EGFP-651For	TCACATGGTCCTGCTGGAGTT
IRES Rev	CTCACATTGCCAAAAGACG
mU6 For	ACAGACTTGTGGGAGAAGC
CMV Rev	GTAGGAAAGTCCCATAAGGTCA
CMV For	GTGGGAGGTCTATATAAGCAGAGCTCG
IRESfor	GTTGTGAGTTGGATAGTTGTGG

Please contact us with any questions:

Viral Vector Core
vectors@uiowa.edu
University of Iowa
500 Newton Road
221 Eckstein Medical Research Building
Iowa City, IA 52242
Tel: (319) 335-6726
<http://www.medicine.uiowa.edu/vectorcore/>

*Updated 2-22-17 AN